

SINGULARITY FORMATION AND COLLAPSE IN THE ATTRACTIVE GROSS-PITAEVSKII EQUATION

A.V. Rybin[†], G.G. Varzugin^{*} and J. Timonen[†]

[†]*Department of Physics, University of Jyväskylä
PO Box 35, FIN-40351 Jyväskylä, Finland*

^{*}*Institute of Physics St. Petersburg State University
198904, St. Petersburg, Russia*

A generic mechanism of collapse in the Gross-Pitaevskii equation with attractive interparticle interactions is gained by reformulating this equation as Newton's equation of motion for a system of particles with a constraint. 'Quantum pressure' effects give rise to formation of a potential barrier around the emerging singularity, which prevents a fraction of the particles from falling into the singularity. For reasonable initial widths of the condensate, the fraction of collapsing particles, which are thereby removed from the condensate, is found to be a 'universal' number $\simeq 0.7$.

The Bose-Einstein condensate formed in a magneto-optically trapped vapor of cold alkali atoms, can be quite accurately described by the Gross-Pitaevskii (GP) equation [1]

$$i\hbar\Psi_t + \frac{\hbar^2}{2m}\Delta\Psi - g\Psi|\Psi|^2 - V(\mathbf{x})\Psi = 0, \quad (1)$$

where $\Psi(\mathbf{x}, t)$ is the wave function of the condensate, the external potential $V(\mathbf{x})$ models the wall-less confinement (the trap), m is the mass of an individual atom, $g = 4\pi\hbar^2 m^{-1}a_s$ is the effective interaction strength (the 'coupling constant') with a_s the scattering length, and $\Delta = \sum_i \frac{\partial^2}{\partial x_i^2}$ is the Laplace operator. A convenient as well as practical choice for the confining trap is the paraboloidal potential $V = \frac{m}{2} \sum_i \omega_i^2 x_i^2$.

It has been known [1–7] for some while already that the Bose-Einstein condensate of atoms with a negative scattering length becomes unstable when the number of atoms exceeds some critical value N_c . These instabilities have recently been analyzed [8] experimentally. To this end the most interesting prospects are provided by the application [9] of Feshbach resonance by which one can actually tune the effective scattering length with an external magnetic field, and thereby e.g. sweep the coupling constant from positive to negative. This technique was very recently used [8] on a condensed gas of ^{85}Rb atoms such that a collapse and the consecutive 'explosion' of a condensate of ^{85}Rb atoms was created.

Triggered by these recent experiments, a possible mechanism through elastic collisions for the ejection from the condensate of atoms with negative scattering length was discussed in [10]. In this Letter we show however

that the mean-field theory (i.e. the GP equation) of the condensate already allows for a description of 'partial collapse', and leads to an accurate estimate for the number of particles that remain in the condensate after the explosion. The fraction of the condensate thrown away in the collapse (i.e. the collapsing part of the condensate), is always about 70% provided that the width of the condensate before the change of sign of the scattering length is close to that of the ground state. This was true e.g. for the experiment reported in [9]. We find that the collapsing fraction is relatively insensitive to the total number of particles in the condensate, and that this fraction tends to 100% only in the limit of very large width of the condensate.

Collapse phenomena are very well known in nonlinear optics and plasma physics, and they have there a long history extending over the last thirty years (see e.g. review [11] and references therein). Following this tradition, the collapse in the GP system with a negative scattering length was discussed in a similar fashion in [12–14].

Within this tradition, the analysis of singularity formation is mainly focussed on the self-similar behaviour of the solutions in the vicinity of the singularity. We have recently shown [15], however, that a true self-similar solution of the GP system is only possible in two space dimensions. This means that a self-similar description of collapse can only be used with certain reservations. In this Letter we propose a completely different description for the singularity formation, which does not rely on self-similarity of the solution. In a sense the starting point of our analysis is the concept of 'partial collapse', now of the Bose-Einstein condensate. A suggestion about a singularity restricted to a localized domain inside the condensate, was already made in [16]. Here we elaborate further on this suggestion, and formulate it within a completely different framework from that used in [16].

Our approach is based on the realization that the GP equation Eq. (1) can be interpreted as Newton's equations of motion for (fictitious) particles with a certain constraint. The dynamics of GP collapse can then easily be followed by following the trajectories of these particles, under the influence of an effective potential due to 'pressure' effect that arise from the spatially varying condensate density. By introducing an auxiliary field $\boldsymbol{\eta}(\mathbf{x}, t)$, and its inverse $\mathbf{r}(\boldsymbol{\eta}, t)$, we can show (see below) that these trajectories satisfy a system of ordinary differential equations,

$$\frac{\partial r_i}{\partial t} = \frac{1}{m} \frac{\partial \phi}{\partial x_i}(\mathbf{r}(\boldsymbol{\eta}, t), t); \quad r_i(\boldsymbol{\eta}, 0) = \eta_i, \quad (2)$$

with $i = 1, 2, 3$ for $d=3$ dimensions. Here ϕ is a phase defined by the representation $\Psi = \sqrt{\rho}e^{i\phi/\hbar}$ of the condensate field. The fields $r_i(\boldsymbol{\eta}, t)$ can now be interpreted as trajectories of some particles with initial points η_i . We describe in the following how the GP equation Eq. (1) can be transformed into the equations of motion for this system of particles.

It is plain that, by using $\Psi = \sqrt{\rho}e^{i\phi/\hbar}$, Eq. (1) is equivalent to a coupled set of equations for ρ and ϕ ,

$$\rho_t + \frac{1}{m} \text{div}(\rho \nabla \phi) = 0 \quad (3)$$

$$\phi_t + \frac{1}{2m}(\nabla \phi)^2 + V + g\rho - \frac{\hbar^2}{2m}\rho^{-1/2}\Delta\rho^{1/2} = 0. \quad (4)$$

Let us now introduce an auxiliary field $\boldsymbol{\eta}(\mathbf{x}, t)$, which is assumed to satisfy

$$\frac{\partial \eta_i}{\partial t} + \frac{1}{m} \nabla \eta_i \nabla \phi = 0. \quad (5)$$

If we define a Jacobian

$$J(\mathbf{x}, t) = \det J_n^m, \quad J_n^m = \frac{\partial \eta_m}{\partial x_n},$$

it then follows from Eq. (5) that

$$\frac{\partial \ln J}{\partial t} + \frac{1}{m} \nabla \ln J \nabla \phi + \frac{1}{m} \Delta \phi = 0. \quad (6)$$

From this Eq. (6) we can easily deduce that

$$\rho(\mathbf{x}, t) = J(\mathbf{x}, t) \rho_0(\boldsymbol{\eta}) \quad (7)$$

is a solution of the continuity equation Eq. (3) with ρ_0 an arbitrary function. We now define the field $\mathbf{r}(\boldsymbol{\eta}, t)$ by requiring that the function $\mathbf{r}(\boldsymbol{\eta}, t)$ is the inverse of the function $\boldsymbol{\eta}(\mathbf{x}, t)$. With this definition Eq. (5) and Eq. (2) become equivalent. Notice that the relation Eq. (7) provides the anticipated representation of the condensate density in terms of particle trajectories.

Thus far we have not used Eq. (4). By differentiating Eq. (2) with respect to t , we find that

$$m \frac{\partial^2 r_i}{\partial t^2} = \frac{\partial}{\partial x_i} \left(\phi_t + \frac{1}{2m} (\nabla \phi)^2 \right). \quad (8)$$

Using now Eq. (4), this Eq. (8) can be expressed in the familiar form of Newton's equations of motion for a set of particles,

$$m \frac{\partial^2 r_i}{\partial t^2} = -\frac{\partial V}{\partial x_i} - \frac{\partial U}{\partial x_i}, \quad (9)$$

in which the 'pressure potential' $U(\mathbf{x}, t)$ is

$$U(\mathbf{x}, t) = -\frac{\hbar^2}{2m} \rho^{-1/2} \Delta \rho^{1/2} + g\rho. \quad (10)$$

The 'particles' of this system thus feel, in addition to the confining trap potential, an effective potential that arises from the spatially and temporally varying condensate density $\rho(\mathbf{x}, t)$. The second term in the right side of Eq. (10) can be recognized as the conventional pressure term, while the first term there arises from the kinetic energy part of the GP equation, and can be considered as a quantum pressure term. As will be explained below, it is the interplay between these two effects in the pressure potential which determines the dynamics of the collapse.

As can be readily seen, the result Eq. (7) can also be written in the form

$$\det \left(\frac{\partial r_i}{\partial \eta_k} \right) \rho(\mathbf{r}, t) = \rho_0(\boldsymbol{\eta}). \quad (11)$$

This constraint can now be used to exclude the particle density ρ from Eq. (9). This means that the system of equations Eqs. (9-11) is self consistent, and fully describes the dynamics of the condensate. We need however to specify first the initial conditions for this system, and a natural choice for Eq. (9) is

$$r_i(\boldsymbol{\eta}, 0) = \eta_i, \quad \frac{\partial r_i}{\partial t}(\boldsymbol{\eta}, 0) = v_i(\boldsymbol{\eta}), \quad (12)$$

with $\mathbf{v} = \frac{1}{m} \nabla \phi|_{t=0}$. The function ρ_0 can now be interpreted as the initial density of the particles, and the phase ϕ can be solved from Eq. (2) up to a time-dependent factor.

Notice that there is a relation between ϕ and ρ . Indeed, from Eq. (2) we find that

$$\det \left(\frac{\partial r_i}{\partial \eta_k} \right) = e^{\frac{1}{m} \int_0^t \Delta \phi(\mathbf{r}, t') dt'}.$$

Hence, we can express ρ in the form

$$\rho(\mathbf{x}, t) = e^{-\frac{1}{m} \int_0^t \Delta \phi(\mathbf{x}(t, t'), t') dt'} \rho_0(\boldsymbol{\eta}(\mathbf{x}, t)), \quad (13)$$

where $\mathbf{x}(t, t') = \mathbf{r}(\boldsymbol{\eta}(\mathbf{x}, t), t')$.

It is evident that a possible collapse of the condensate is related to the singular points of the solution, i.e. to the points where the function $\mathbf{r}(\boldsymbol{\eta}, t)$ is not invertible and the determinant $\det(\frac{\partial r_i}{\partial \eta_k})$ vanishes. At these points the density becomes infinite. It seems natural to assume that if collapse takes place, it happens at a finite time $t = t_*$, i.e. $\rho(\mathbf{r}_s, t_*) = \infty$. Then, from Eq. (11), we can conclude that for any $\mathbf{r}_s$ there is a set Ω_s such that

$$\lim_{t \rightarrow t_*} r_i(\boldsymbol{\eta}, t) = r_{si}$$

for all $\boldsymbol{\eta} \in \Omega_s$. In other words, all particle trajectories that begin at any $\boldsymbol{\eta} \in \Omega_s$ are converged at one point $\mathbf{r}_s$.

Suppose that the set of singular points, $\{\mathbf{r}_s\}$, consists of one isolated point while Ω_s is a domain. We can then define the number of particles involved in the collapse as

$$N_e = \int_{\Omega_s} \rho_0(\boldsymbol{\eta}) d^3\boldsymbol{\eta}. \quad (14)$$

This number can thus be interpreted as the number of particles that are ejected from the condensate when it is collapsed. Notice that the domain Ω_s can also coincide with the whole space. In this case $N_e = N$, the total number of atoms in the condensate. This is true in particular for the self-similar solutions, when the whole condensate collapses to a δ -function centered in the trap, and completely disappears apart from the very particular borderline case of an oscillating solution [15]. The self-similar solutions can only exist [15] in two space dimensions. It is important to emphasize that, *in general*, $N_e < N$. It can be shown, *e.g.*, that if ρ_0 tends to a Gaussian distribution as $|\boldsymbol{\eta}| \rightarrow \infty$, then $N_e < N$. This means that N_e is an essential characteristics of the collapse of attractive condensates.

We can explain the partial collapse qualitatively as follows. If, initially, the width of the condensate is greater than the characteristic oscillator length a_{HO} , both the quantum pressure and the conventional pressure induce the condensate to decrease its width. However, when the width becomes equal or less than a_{HO} , the two pressure terms in the energy have opposite effects: quantum pressure favours an expanding condensate (as a result of an 'uncertainty principle'), while the negative conventional pressure term favours a collapsing condensate. Provided that the number of particles is greater than a critical value, the negative pressure dominates over the quantum pressure in a small domain centered at a point $\mathbf{r}_s$. Outside this domain it is the quantum pressure that dominates, and prevents the outside particles from entering this domain.

In order to estimate the value of N_e , we use here a Gaussian trial wave function [17–20]. It gives an approximate solution for the density, which can be expressed in the form

$$\rho(\mathbf{x}, t) = \frac{N}{\pi^{3/2} \prod_i \tau_i(t)} e^{-\sum_i \frac{x_i^2}{\tau_i^2(t)}}, \quad (15)$$

in which the functions τ_i satisfy [21–23]

$$\ddot{\tau}_i + \omega_i^2 \tau_i - \frac{\hbar^2}{m^2} \frac{1}{\tau_i^3} - \frac{gN}{(2\pi)^{3/2} m} \frac{1}{\prod_k \tau_k} \frac{1}{\tau_i} = 0 \quad (16)$$

with $\tau_i(0) = a_i$ and $\dot{\tau}(0) = 0$. Here a_i is the initial width of the condensate.

Substituting the solution Eq. (15) in Eq. (9), we find that

$$\begin{aligned} \frac{\partial^2 r_i}{\partial t^2} &= -\frac{\partial V_{eff}}{\partial r_i} \\ V_{eff}(\mathbf{r}, t) &= \frac{1}{2} \sum_i \left(\omega_i^2 - \frac{\hbar^2}{m^2 \tau_i^4} \right) r_i^2 + \frac{g}{m} \rho(\mathbf{r}, t). \end{aligned} \quad (17)$$

Here V_{eff} is an effective time-dependent potential that determines the behaviour of particle trajectories in the vicinity of the center of the trap.

For simplicity, we consider here in more detail the symmetric trap: $\omega_i = \omega$ and $a_i = a$. Hereafter we measure the time in units of $1/\omega$, and the distance in units of $a_{HO} = \sqrt{\hbar/m\omega}$, the oscillator length. Now Eqs. (17),(16) take the form

$$\begin{aligned} \frac{d^2 r}{dt^2} &= -\frac{\partial V_{eff}}{\partial r} \\ V_{eff} &= \frac{1}{2} \left(1 - \frac{1}{\tau^4} \right) r^2 - \frac{4}{\sqrt{\pi}} \frac{g_0}{\tau^3} \left(e^{-\frac{r^2}{\tau^2}} - 1 \right), \end{aligned} \quad (18)$$

and

$$\frac{d^2 \tau}{dt^2} + \tau - \frac{1}{\tau^3} + \sqrt{\frac{2}{\pi}} \frac{g_0}{\tau^4} = 0, \quad (19)$$

in which $g_0 = \frac{|a_s|N}{a_{HO}}$, and the initial conditions are given by $r(0) = \eta$, $\dot{r}(0) = 0$, $\tau(0) = a$, $\dot{\tau}(0) = 0$. Notice that if $g_0 > g_{0cr}$, where $g_{0cr} \approx 0.671$ [1], the solution of Eq. (19) has a zero at a finite time $t = t_*$.




FIG. 1. The effective potential V_{eff} at different times $t < t_*$. The initially almost paraboloidal potential develops a growing barrier at the final stages of the collapse.

Let us assume that the system has $a > 1$ and begins to collapse. Initially, V_{eff} can be well approximated by a paraboloidal trap. The particles are drifted towards the center and the condensate width decreases. When this width is reduced to the oscillator length, a potential barrier is however formed at a distance $r = r_{max}$ from the center,

$$r_{max} = \tau \sqrt{\ln \frac{8g_0}{\sqrt{\pi}(\tau - \tau^5)}},$$

preventing the outside particles from falling into the emerging singularity. Figure 1 illustrates the formation

of this potential barrier in the limit when the width of the condensate τ goes to zero.

We solved the system Eqs. (18),(19) numerically. The collapse time was obtained from the condition¹ $\tau(t_*) < 0.1$. Equation (18) was solved in the time interval $[0, t_*]$, while the point η was supposed to belong to the domain Ω_s provided that $r(t_*) < r_{max}(t_*)$.

The conditions of the experiment reported in [9] were modelled such that the initial width of the condensate was derived as a stationary solution of Eq.(19) with g_0 replaced by $-41.67g_0$. Notice that in the experiment [9] the initial positive scattering length was $2500a_0$ (a_0 is the Bohr radius), and this scattering length was in 0.5ms changed to $-60a_0$. The number -41.67 is thus the ratio of these two scattering lengths. We varied the parameter g_0 in a large range, from 1 to 300. In all cases the fraction of the particles ejected from the condensate as a result of collapse, was about 70%. We also found that the fraction of ejected particles increased above 70% only when the initial width of the condensate was increased well above that of the ground state. In the limit of very large widths the fraction approached 100% of the total number of particles.

In summary, in this Letter we have explained the generic mechanism of collapse in a trapped Bose-Einstein condensate with attractive interparticle interactions in the framework of mean-field theory (GP equation). We can conclude that, in general, only a fraction of the condensate takes part in the collapse, while the rest of the particles are screened from the emerging singularity by an effective potential barrier due to 'quantum pressure' effects. This potential barrier is somewhat reminiscent of the 'event horizon' that appears around gravitational singularities. We could also show that, for reasonable initial widths of the condensate, the fraction of particles removed from the condensate as a result of collapse is an almost 'universal' number $\simeq 0.7$. Even though we here derived this fraction of ejected particles only approximately, our definition of the related particle number N_e Eq. (14), can be made mathematically rigorous. This number can be obtained from the solution of Eq. (9). The simplified model based on Gaussian trial wave function, which we proposed here, is not accurate if the 'coupling constant' g_0 is in the vicinity of g_{0cr} . A more thorough analysis of Eq. (9) will be reported in a forthcoming publication.

¹The final stage of the collapse takes place very fast, so a good numerical estimate for t_* was already obtained from this seemingly rather crude approximation, which however allowed us to avoid the difficulties related to the mathematical singularity at $t = t_*$.

ACKNOWLEDGMENTS

This work was supported by the Academy of Finland under the Finnish Center of Excellence Programme 2000-2005 (Project No. 44875, Nuclear and Condensed Matter Physics at JYFL). GGV was partly supported by an RFBR grant No. 00-01-00480.

-
- [1] F. Dalfovo, S. Giorgini, L.P. Pitaevskii, and S. Stringari, Rev. Mod. Phys. **71**, 463 (1999).
 - [2] F. Dalfovo, and S. Stringari, Phys. Rev. A, **53**, 2447 (1996).
 - [3] R.J. Dodd, K. Burnett, Mark Edwards, and C.W. Clark, Phys. Rev. A **56**, 587 (1997).
 - [4] C.C. Bradley, C.A. Sackett, J.J. Tollet, and R.G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).
 - [5] C.A. Sackett, C.C. Bradley, M. Welling, and R.G. Hulet, Appl. Phys. B: Lasers Opt. **65**, 433 (1997).
 - [6] M. Houbiers, and H.T.C. Stoof, Phys.Rev. A **54**, 5055 (1996).
 - [7] Yu. Kagan, G.V. Shlyapnikov, and J.T.M. Walraven, Phys. Rev. Lett. **76**, 2670 (1996).
 - [8] C. A. Sackett, J. M. Gerton, M. Welling and R. G. Hulet, Phys. Rev. Lett. **82**, 876 (1999).
 - [9] S. L. Cornish, N. R. Claussen, J. L. Roberts, E. A. Cornell and C. E. Wieman, Phys. Rev. Lett. **85**, 1795 (2000).
 - [10] R. A. Duine and H. T. C. Stoof, cond-mat/0007320 (2000).
 - [11] J. J. Rasmussen and K. Rypdal, Physica Scripta **33**, 481 (1986); K. Rypdal and J. J. Rasmussen Physica Scripta **33**, 498 (1986).
 - [12] L. P. Pitaevskii, Phys. Lett. A **221**, 14 (1996).
 - [13] M. Wadati and T. Tsurumi, Phys. Lett. A **247**, 287 (1998).
 - [14] L. Bergé, T.J. Alexander, and Y.S. Kivshar, Phys. Rev. A **62**, 023607 (2000).
 - [15] A. V. Rybin, G. G. Varzugin, M. Lindberg, J. Timonen and R. K. Bullough, Phys. Rev. E **62**, 6224 (2000).
 - [16] M. Ueda and K. Huang, Phys. Rev. A, **60**, 3317 (2000).
 - [17] G. Baym, and C. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
 - [18] H.T.C. Stoof, J. Stat. Phys. **87**, 1353 (1997).
 - [19] A. Parola, L. Salashnich, and L. Reatto, Phys. Rev. A **57**, R3180 (1998).
 - [20] A.L. Fetter, J. Low Temp. Phys. **106**, 643 (1997).
 - [21] V.M. Perez-Garcia, H. Michinel, J.I. Cirac, M. Lewenstein, and P. Zoller, Phys.Rev.Lett. **77**, 5320 (1996); Phys.Rev.A **56**, 1424 (1997).
 - [22] Yu. Kagan, E.L. Surkov, and G.V. Shlyapnikov, Phys. Rev. Lett. **55**, R18 (1997).
 - [23] F. Dalfovo, C. Minniti, S. Stringari, and L. Pitaevskii, Phys. Lett. A **227**, 259 (1997).